

BIOMIN® POTASSIUM 2-0-25



Mineral for Soil and Foliar Application

GENERAL INFORMATION

BiomIn Potassium 2-0-25 is a liquid plant food designed for soil and/or foliar applications. **BiomIn Potassium** is compatible with most fertilizer solutions and pesticides. Dilute **BiomIn Potassium** with water before mixing with other fertilizers or pesticides.

DIRECTIONS OF USE:

Shake well before using.

SOIL APPLICATION:

BiomIn Potassium can be applied before or at planting as banding or broadcast. The application rate is 2-3 gallons per acre with at least 50 gallons of water.

FOLIAR APPLICATION:

Dilute **BiomIn Potassium** with at least 50 times of water and apply to plants according to the following recommendations:

FIELD CROPS: Apply 2-4 quarts per acre of **BiomIn Potassium** when crops are about 4-6 inches (10-20 cm) high or when deficiency symptoms appear. Repeat the application in 2 - 4 weeks if necessary.

Alfalfa - 2-4 quarts per acre one week after each cutting.

Beans, Peas - 2-4 quarts per acre at bud stage, prior to bloom. Repeat the application 2 more times in 2 week intervals.

Citrus - 2-4 quarts per acre at bud stage, prior to bloom. When new growth appears, repeat the application 2 times at 2-4 week intervals.

GUARANTEED ANALYSIS

Total Nitrogen(N)2%
2% Water Soluble Organic Nitrogen

Soluble Potash (K_2O)25%

Contains Potassium Carbonate, Citric Acid and Glycine as Complexing Agent. (Nitrogen derived from Glycine.)

ALSO CONTAINS NON PLANT FOOD INGREDIENTS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY AT TIME OF BLOOM

Manufactured by:
JH Biotech, Inc.
P.O. Box 3538 Ventura, CA 93006

Net Contents:

- 2.5 Gallons (9.45 liters)
- 2 x 2.5 Gallons (18.9 liters)
- 5 Gallons (18.9 liters)
- 55 Gallons (207.9 liters)
- Others: _____ Gallons

Lot No.:

Cotton - 2-4 quarts per acre after true leaves appear. Second application should be made at early square at 2-3 quarts per acre. Repeat the application 2 more times at 2 week intervals.

Potato - 2-4 quarts per acre at 6-8 leaf stage. 2-4 quarts per acre at tuber initiation. 2-4 quarts per acre when tubers are 1"-2" in diameter. Repeat the application 2 more time in 1-2 week intervals.

Strawberry - 2-4 quarts starting after plants are established. Repeat application at 2-3 week intervals throughout the season.

Tomato and Pepper - 2-4 quarts per acre after stand is established. 2-4 quarts per acre at bud stage, prior to bloom. Two additional applications at 2-4 quarts per acre at 2 week intervals.

BiomIn Potassium IS RECOMMENDED FOR USE ON:

Almond, Apple, Apricot, Avocado, Barley, Beans, Carrot, Corn, Grapes, Hops, Peach, Pear, Pecan, Pistachio, Plum, Rice, Sorghum, Soybean, Sugarbeet, Vegetables, Walnut

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aspfco.org/metals.htm>.

JH Biotech, Inc.

0836-D-A-1
Copyright © 2006 JH Biotech, Inc.
All rights reserved.