

BIOMIN[®] MAGNESIUM POWDER

5-0-0

Chelated Mineral for Soil and Foliar Application



GUARANTEED ANALYSIS

Total Nitrogen (N).....	5.0%
5% water soluble organic nitrogen	
Magnesium (Mg).....	12.0%
12% chelated magnesium	

Derived from magnesium sulfate and citric acid. Nitrogen derived from glycine.

PRODUCT INFORMATION

BIOMIN MAGNESIUM contains nitrogen and chelated magnesium for soil and/or foliar applications. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. **BIOMIN MAGNESIUM** is designed to prevent and correct magnesium deficiency and to boost crops during critical or fast growing periods. May be applied with nitrogen fertilizers.

BIOMIN MAGNESIUM can be applied to ALL crops including all vegetables, potatoes, carrots, onions, all berries and strawberries, all brassicas, all fruit trees, avocados, persimmons, citrus, all nut trees, vineyards, kiwifruit, tropical fruit, lettuce, asparagus, all melons, beans, ginger, ornamentals, cut flowers and pastures.

CAUTION - Do not apply this product after application of copper containing products as this may cause phytotoxicity.

KEEP OUT OF REACH OF CHILDREN

PREVENTS AND CORRECTS MINERAL DEFICIENCIES

COMPATIBILITY: Always perform a compatibility test before mixing any products with **BIOMIN MAGNESIUM**.

DIRECTIONS AND GENERAL RECOMMENDATIONS:

SOIL APPLICATION

BIOMIN MAGNESIUM may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1.5 - 5.0 LBS/Acre (1.0-5.5 Kg/Ha) depending on crop, soil conditions and application methods. For maintenance, 1.5 LBS/Acre (1.5 Kg/Ha) are usually recommended.

FOLIAR APPLICATION

FIELD CROPS: Apply 1.0-1.5 LBS/Acre (1.0-1.5 Kg/Ha) of **BIOMIN**

MAGNESIUM when crops are 6-12 inches (15-30 cm) high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

VEGETABLES and POTATOES: Apply 1.0-1.5 LBS/Acre (1.0-1.5 Kg/Ha) of **BIOMIN MAGNESIUM** at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later.

GREENHOUSE CROPS: Apply 1.0-1.5 LBS/Acre (1.0-1.5 Kg/Ha) of **BIOMIN MAGNESIUM** 2-3 weeks after emergence or transplanting, or when magnesium deficiencies first appear. Repeat the application every one or two weeks.

FRUIT TREES AND GRAPES: Apply 1.0-1.5 LBS/Acre (1.0-1.5 Kg/Ha) of **BIOMIN MAGNESIUM** in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

BIOMIN MAGNESIUM is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 200 times of water. For best results and more specific recommendations, contact your nearest distributor.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538, Ventura, CA 93006, USA

NET CONTENTS: 25LB (11.36KG) 50 LB (22.72KG)

U.S. Pat.# 5,504,055

LOT NO.:

JH Biotech, Inc.

0404-CA-G-2
Copyright © 2011 JH Biotech, Inc.
All rights reserved.