

BUFFERMIN® - L

Glycine Chelated Minerals



Product Information

BUFFERMIN-L contains glycine chelated minerals intended to be used as a nutritional supplement for animals. It can be used in drinking water, mild, and milk replacer.

GUARANTEED ANALYSIS

Calcium (Ca).....1,000 ppm
Magnesium (Mg).....10,000 ppm
Cobalt (Co).....24 ppm
Copper (Cu).....220 ppm
Iron (Fe).....520 ppm
Manganese (Mn).....1,100 ppm
Zinc (Zn).....3,200 ppm

***Density:** 1.06 g/ml

INGREDIENTS:

Calcium glycinate, cobalt glycinate, copper glycinate, iron glycinate, magnesium glycinate, manganese glycinate and zinc glycinate.

KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USE

NET CONTENTS: 20 liters

Manufacturing Date:

Expiration Date:

Lot #:

Directions and General Recommendations

FEEDING INSTRUCTIONS:

BUFFERMIN-L maybe used in drinking water, milk or milk replacers.

RECOMMENDED RATES ARE:

Chicks and Poults:	Mix 2 milliliter of Buffermin - L per liter of drinking water for the first 10 days after hatching.
Pullets:	Mix 2 milliliter of Buffermin - L per liter of drinking water for the first 10 days after entering laying house.
Hens:	Mix 2 milliliter of Buffermin - L per liter of drinking water.
Baby pigs:	Dilute 5 milliliter of Buffermin - L with 100 milliliter of milk and give 1 – 2 milliliter to each baby pig per day.
Weaning pigs:	Mix 2 milliliter of Buffermin - L per liter of drinking water for 10 days.
Feeder pigs:	Mix 2 milliliter of Buffermin - L per liter of drinking water during and after transportation and continue for 10 days.
Sows:	Mix 2 milliliter of Buffermin - L per liter of drinking water 2 weeks prior to and after farrowing.
Dairy cows:	Mix 2 milliliter of Buffermin - L per liter of drinking water.
Calves:	Mix 15 – 30 milliliters of Buffermin - L in 60 milliliter of milk to administer as a drench. Mix 15 milliliter of Buffermin – L per head per day in milk replacer.
Feeder cattle:	Mix 2 milliliters of Buffermin - L per liter of drinking water for 5 – 10 days.
Lambs:	Mix 2 milliliters of Buffermin - L per liter of drinking water.
Horses:	Mix 2 milliliters of Buffermin - L per liter of drinking water.

Please Store in dry and cool places.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

JH Biotech Inc.

1301-I-D-1

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538, Ventura, CA 93006 U.S.A. Phone: 805-650-8933 Website: www.jhbiotech.com

Copyright © 2013 All rights reserved